



MISSOURI CARE BEHAVIORAL HEALTH INFORMATION



TREATMENT OF ALCOHOL AND OTHER DRUG DEPENDENCE

We have adopted the National Institution Alcohol Abuse and Alcoholism (NIAAA), Helping Patients Who Drink Too Much, A Clinician's Guide, 2005 Edition. You can find the text of this guideline as well as other tools online at: http://pubs.niaaa.nih.gov/publications/Practitioner/CliniciansGuide2005/clinicians_guide.htm.

As an addendum to the NIAAA Helping Patients Who Drink Too Much, A Clinician's Guide, 2005 Edition, the following guidelines for initiation and engagement of alcohol and other drug dependence treatment have been adopted based on the Healthcare Effectiveness Data and Information Set, (HEDIS):

- **Initiation of Alcohol and Other Drug (AOD) Dependence Treatment**
Adolescent and adult members diagnosed with AOD dependence should initiate treatment

through either an inpatient AOD admission or an outpatient service for AOD dependence and additional AOD services within 14 days.

- **Engagement of Alcohol and Other Drug (AOD) Dependence Treatment**

Adolescent and adult members diagnosed with AOD dependence should engage in treatment with 2 additional AOD services 30 days after initiation.

No guideline should be considered the sole source of guidance for the treatment of a particular disorder. They are designed to provide a framework for treatment decisions. They are not intended as a protocol of treatment measures or to replace the clinical judgment of the provider.

Missouri Care, an Aetna Health Plan

For additional information, please contact Ronald L Lacey, MD,
Medical Director of Behavioral Health at ron.lacey@aetna.com or at 800-889-4073.

Questions?

Call Behavioral Health Services
at 1-800-889-4073
www.MissouriCare.com



Missouri CareSM
AN AETNA HEALTH PLAN